

100 pts. Show your work. If you use your NSpire for a computation, make a note of that fact.

1. Consider the differential equation $y'' + 2y' + y = 0$

a. What is the order of the equation?

b. Verify that $y(x) = e^{-x}$ is a solution of the equation

c. What is the phrase that describes a DE together with an Initial Condition?

2. Consider the first order linear DE $y' + \frac{3}{x}y = \frac{e^x}{x^3}$.


a. Use the integrating factor to find a general solution to the equation

b. Use the general solution to explain briefly why all of the solutions have as an asymptote the negative x-axis

3. Consider the separable first-order DE $y' = x^2\sqrt{y}$.

a. Find a general solution

b. Sketch the slope field, setting your window as shown below (two zoom in's at (0,1) will be close).




c. Explain briefly why, looking at the **general solution**, that the solutions are defined only in quadrants I and II.

d. Add to the graph the solutions that correspond to IC's $y(0) = 2$, $y(0) = 1$, and $y(0) = 0$.


e. Write a different IC (approximate) that will generate the same solution as $y(0) = 2$

4. Match the slope field to the differential equation. On each curve, sketch the solution that corresponds to $y(1)=-1$. Then give a second IC that generates the same particular solution. One “zoom in” at $(0,0)$ will be close to my window settings.


- a. $y' = \sin(x) + \sin(y)$ b. $y' = e^{-\frac{x}{2}} + 2y$ c. $y' = 0.2x - \sin y$ d. $y' = \sqrt[3]{x} - y - 2$




Eq: _____ Other IC: _____



Eq: _____ Other IC: _____



Eq: _____ Other IC: _____



Eq: _____ Other IC: _____


5. Consider the DE $(y + 2xy^3)dx + (1 + 3x^2y^2 + x)dy = 0$

a. Verify that the equation is exact

b. Find the general solution

c. Find the particular solution for $y(1) = -1$

d. Sketch the slope field and the particular solution from part c.



6. Consider the 1st order linear differential equation $y' + 3x^2y = xe^{-x^3}$.

a. Write the complementary homogeneous equation.

b. Solve the complementary equation and label the solution y_1 .

c. Find the general solution y_g to the equation using variation of parameters.

7. For **any two of these three equations**, find a general solution. Choose your own method, **show your work**.

a. $(x + \sin y) + (x \cos y - 2y)y' = 0$

b. $xy' + y = x \cos x$

c. $y' = \sqrt[3]{8xy}$

8. A missile is launched straight upward from a silo positioned 80 ft below ground level, with initial velocity 500 ft/sec. Assume that the acceleration due to gravity is 32 ft/sec/sec. Use two decimal places where needed.
- Find formulas for the acceleration, velocity, and altitude of the projectile
 - Find the maximum height it achieves
 - After how many seconds does it leave the silo (cross ground level on the way up)?
 - What is the velocity when it leaves the silo?

Extra Credit: Prove that any separable equation is exact.

Formulas which you may or may not use on this exam

$c < y < d$. Then the differential equation

$$M(x, y) dx + N(x, y) dy = 0$$

is exact in R if and only if

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

- a.
- b. For a first order linear differential equation $y' + p(x)y = f(x)$,
- To use the integrating factor, $y_g = \frac{1}{\rho} \times \int \rho \cdot f(x) dx$
 - To use variation of parameters, $y_1 = e^{-\int p(x) dx}$, $u = \int \frac{f(x)}{y_1(x)} dx$, and $y_g = u \cdot y_1$
- c. Motion relationships:
- $v(t) = \int a(t) dt$
 - $s(t) = \int v(t) dt$
 - For vertical projectile motion with gravity the only acceleration, $h(t) = -\frac{g}{2} t^2 + v_0 t + h_0$
- d. An equation is **separable** if it can be written $y' = \frac{g(x)}{h(y)}$, or $h(y) dy = g(x) dx$